

Heavy Flavor Production in PHENIX

O. Drapier

LLR-École Polytechnique, France
for the PHENIX Collaboration


✓ "Onia" production

✓ Leading order at low x
= "gluon fusion"

✓ Sensitive to:

✓ Initial state

- ✓ Parton distribution functions
- ✓ p_T broadening
- ✓ Parton energy loss in the initial state ?
- ✓ Polarization ?



✓ Final state

- ✓ Parton energy loss in the hot & dense medium ?
- ✓ Thermal enhancement ?
- ✓ Flow ?

+ feed-down (e.g. B or $\chi_c \rightarrow J/\psi$)


✓ Open charm (or beauty) production

- ✓ Leading order at low x
= "gluon fusion"

✓ Sensitive to:

✓ Initial state

- ✓ Parton distribution functions
- ✓ p_T broadening
- ✓ Parton energy loss in the initial state ?
- ✓ Polarization ?






✓ Final state

- ✓ Parton energy loss in the hot & dense medium ?
- ✓ Thermal enhancement ?
- ✓ Flow ?

- ✓ Open flavors are interesting *per se* :
- ✓ Do heavy quark suffer « quenching », as light flavors do ?
 - ✓ Higher quark mass \rightarrow less gluon radiation ("dead cone effect")
 - ✓ D/π modified at moderate p_T due to different quenching
 - ✓ Y. L. Dokshitzer & D.E. Kharzeev, Phys.Lett.B519(2001)199-206
- ✓ Do heavy quark flow ?
 - ✓ Early creation in hard process \rightarrow they should not
 - ✓ Additional creation, enhancement, "in medium" effects \rightarrow they could
 - ✓ In any case, OPEN = they're detected once bound to light quarks
 - ✓ Influence of light quark flow
 - ✓ Influence of decay if detected in semi-leptonic decay channels






- ✓ Possible charm enhancement in Heavy Ion collisions ?
 - ✓ "unexpected sources" of dileptons observed at SPS
 - ✓ This contribution doesn't scale as the number of collisions






- ✓ Is there an enhancement of charm production ?
- ✓ Ideally, open charm should be used as a reference for J/ ψ production, as color screening prevents charmonia bound states

color screening \Leftrightarrow bound charmonia / open charm

- ✓ Heavy flavor = probe for « cold » nuclear effects
 - ✓ Parton distribution functions are modified in nuclei
 - color screening \Leftrightarrow bound charmonia / open charm**
- ✓ e.g. in d-Au collisions :



- ✓ Charm production is sensitive to incident energy of partons
 - ✓ Possible energy loss in the initial state may affect charmonium production
 - ✓ Holds both for charmonia and open charm, so could be checked with open charm




- ✓ Sensitive p_T broadening
(e.g.: gluon-nucleon scattering)


color screening \Leftrightarrow bound charmonia / open charm

✓ RHIC (Relativistic Heavy Ion Collider)

- ✓ Long Island, not so far from Manhattan ;-)
- ✓ Dedicated to heavy ion physics & spin studies
- ✓ 4 experiments
- ✓ 100+100 GeV/A
- ✓ Variable incident energy
- ✓ p-p up to 500 GeV




- ✓ Vast incident energy range
- ✓ Various ion species from p to Au




- ✓ Electrons in EM-Cal + RICH
- ✓ Muons in MuTr + MuId



- ✓ Secondary decay vertex ... long term future ?
 - ✓ Needs precise tracking close to IP to see displaced vertex ...
 - ✓ we have no detector to do this
 - ✓ Very challenging in high multiplicity & radiation environment
- ✓ D reconstruction ($K\pi$)
 - ✓ The best way ... still under study in PHENIX
- ✓ Semi-leptonic decays
 - ✓ Single leptons (e or μ)
 - ✓ Dilepton continuum (e^+e^- or $\mu^+\mu^-$)
 - ✓ $e-\mu$ coincidences





- ✓ Different acceptance domains



- ✓ For now:
 - ✓ Only single electrons in central arms
 - ✓ Only open charm

- ✓ Electrons
 - ✓ $|\eta| < 0.35$ $\Delta\phi = \pi / 4$
 - ✓ Tracks reconstructed with Drift Chamber & PC1
 - ✓ EM-Cal matching hit
 - ✓ RICH hits + ring shape
 - ✓ Timing (EMC or TOF)

- ✓ "Photonic" electrons
 - ✓ γ conversion
 - ✓ π^0 and η/η' Dalitz decay: $\rightarrow \gamma ee$
 - ✓ light vector meson decay:
 - ✓ $\omega, \phi \rightarrow (\pi^0, \eta) ee$





- ✓ NON-photonic electrons = ALL - PHOTONIC
 - ✓ K decay
 - ✓ $\rho, \omega, \phi \rightarrow ee$
 - ✓ $c \rightarrow e$ (dominant)
 - ✓ $b \rightarrow e$


- ✓ Subtract background by:
 - ✓ Cocktail method
 - ✓ Converter method
 - ✓ Direct measurement of γe coincidences + event mixing

- ✓ Correct for acceptances and efficiencies


- ✓ Non-photonic contribution increases with p_T
 - ✓ Good agreement between cocktail method and γe coincidence method




- ✓ Non-photonic electrons in p-p as compared to Pythia:
 - ✓ Reasonable agreement
 - ✓ Charm alone not sufficient at high p_T
 - ✓ Bottom \rightarrow not enough statistics




- ✓ Phenomenological fit
 - ✓ To compare with d-Au and Au-Au ...




✓ ... and to extract charm cross section



$$\sigma_{cc} = 709 \mu\text{b} \pm 85(\text{stat})^{+332}_{-281}(\text{syst})$$







- ✓ d-Au collisions: minimum bias result (binary scaled)






- ✓ COMPATIBLE WITH BINARY SCALING

✓ d-Au: binary scaling also holds for different centralities



- ✓ E_{ZDC} (spectators) / BBC (secondary particles) correlation
 - ✓ "zero degree calorimeter" + "beam-beam counter"
- ✓ As % of total cross-section










B (fm)	N_{part}	N_{coll}
2,3 $\pm 0,9$	353 ± 19	1091 ± 102
7,1 $\pm 0,5$	181 ± 16	422 ± 65
14,5 $\pm 0,3$	4,1 $\pm 2,5$	2,8 $\pm 2,2$


- ✓ Glauber model
- ✓ Nuclear density profile
- ✓ Geometry
- ✓ N-N cross-section

$1/T_{AB}$ dN/dp^3 [mb GeV $^{-2}$ c 3]

- ✓ Au-Au as a function of centrality
- ✓ Compatible with binary scaling
 - ✓ Need more stat. at high p_T !!!




✓ Au-Au as a function of centrality: summary plot




✓ Open charm in Au-Au : consistent with N_{coll} scaling


- ✓ $0.8 < p_T < 4.0$
- ✓ $\alpha = 0.938 \pm 0.075$ (stat) ± 0.018 (syst)
- ✓ Together with p-p: $\alpha = 0.958 \pm 0.035$ (stat)



- ✓ Open charm in Au-Au @ 62.4 GeV, as compared to ISR p-p data at the same incident energy
 - ✓ Also compatible with $\langle N_{\text{coll}} \rangle$ scaling




- ✓ Undistinguishable scenarios from p_T distributions
 - ✓ Would require high statistics at high p_T
 - ✓ Difference smoothed by decay



S. Batsouli, S.Kelly, M.Gyulassy, J.Nagle
Phys.Lett. B557 (2003) 26-32

- ✓ What about direct flow measurement ... V_2 ?
 - ✓ V_2 of non photonic electrons in 200 GeV/A Au-Au (run 2)
 - ✓ Need more statistics (run 4)




- ✓ PHENIX measures open charm with single electrons
- ✓ Proton-proton : reasonable agreement with PYTHIA
 - ✓ Charm + bottom
 - ✓ σ somewhat lower than reported by STAR
- ✓ d-Au: good agreement with Ncoll scaling for the 4 centrality bins
- ✓ Au-Au : good agreement with Ncoll scaling for the 5 centrality bins: $\alpha = 0.938 \pm 0.075$ (stat) ± 0.018 (syst)
 - ✓ Together with p-p: $\alpha = 0.958 \pm 0.035$ (stat)
- ✓ NO DEVIATION FROM BINARY SCALING OBSERVED SO FAR IN OPEN CHARM PRODUCTION

✓ Near future: Run4 = much more statistics to come !

- ✓ + Single muons
- ✓ + Dilepton continuum
 - ✓ needs high stat + good control of combinatorial background
- ✓ + Electron-muon coincidences ?

✓ PHENIX upgrades:

- ✓ Barrel VTX
 - ✓ Pixel + strips
- ✓ Forward VTX
 - ✓ strips



Future heavy flavor detection in PHENIX:

- Beauty and low p_T charm via displaced e and/or μ $-2.7 < \eta < -1.2, |\eta| < 0.35, 2.7 < \eta < 1.2$
- Beauty through displaced $J/\psi \rightarrow ee$ ($\mu\mu$) $-2.7 < \eta < -1.2, |\eta| < 0.35, 2.7 < \eta < 1.2$
- High p_T charm through $D \rightarrow \pi K$ $|\eta| < 0.35$

Brazil	University of São Paulo, São Paulo
China	Academia Sinica, Taipei, Taiwan China Institute of Atomic Energy, Beijing Peking University, Beijing
France	LPC, University de Clermont-Ferrand, Clermont-Ferrand Dapnia, CEA Saclay, Gif-sur-Yvette IPN-Orsay, Universite Paris Sud, CNRS-IN2P3, Orsay LLR, Ecole Polytechnique, CNRS-IN2P3, Palaiseau SUBATECH, Ecole des Mines at Nantes, Nantes
Germany	University of Münster, Münster
Hungary	Central Research Institute for Physics (KFKI), Budapest Debrecen University, Debrecen Eötvös Loránd University (ELTE), Budapest
India	Banaras Hindu University, Banaras Bhabha Atomic Research Centre, Bombay
Israel	Weizmann Institute, Rehovot
Japan	Center for Nuclear Study, University of Tokyo, Tokyo Hiroshima University, Higashi-Hiroshima KEK, Institute for High Energy Physics, Tsukuba Kyoto University, Kyoto Nagasaki Institute of Applied Science, Nagasaki RIKEN, Institute for Physical and Chemical Research, Wako RIKEN-BNL Research Center, Upton, NY Rikkyo University, Tokyo, Japan Tokyo Institute of Technology, Tokyo University of Tsukuba, Tsukuba Waseda University, Tokyo
S. Korea	Cyclotron Application Laboratory, KAERI, Seoul Kangnung National University, Kangnung Korea University, Seoul Myong Ji University, Yongin City System Electronics Laboratory, Seoul Nat. University, Seoul Yonsei University, Seoul
Russia	Institute of High Energy Physics, Protovino Joint Institute for Nuclear Research, Dubna Kurchatov Institute, Moscow PNPI, St. Petersburg Nuclear Physics Institute, St. Petersburg St. Petersburg State Technical University, St. Petersburg
Sweden	Lund University, Lund



12 Countries; 58 Institutions; 480 Participants*

USA	Abilene Christian University, Abilene, TX Brookhaven National Laboratory, Upton, NY University of California - Riverside, Riverside, CA University of Colorado, Boulder, CO Columbia University, Nevis Laboratories, Irvington, NY Florida State University, Tallahassee, FL Florida Technical University, Melbourne, FL Georgia State University, Atlanta, GA University of Illinois Urbana Champaign, Urbana-Champaign, IL Iowa State University and Ames Laboratory, Ames, IA Los Alamos National Laboratory, Los Alamos, NM Lawrence Livermore National Laboratory, Livermore, CA University of New Mexico, Albuquerque, NM New Mexico State University, Las Cruces, NM Dept. of Chemistry, Stony Brook Univ., Stony Brook, NY Dept. Phys. and Astronomy, Stony Brook Univ., Stony Brook, NY Oak Ridge National Laboratory, Oak Ridge, TN University of Tennessee, Knoxville, TN Vanderbilt University, Nashville, TN
-----	---

*as of January 2004